

q -VOLKENBORN INTEGRATION AND ITS APPLICATIONS

TAEKYUN KIM

Department of Mathematics Educations, Kongju
National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr

ABSTRACT. The main purpose of this paper is to present a systemic study of some families of multiple q -Euler numbers and polynomials. In particular, by using the q -Volkenborn integration on $\mathbb{Z}_p$, we construct p -adic q -Euler numbers and polynomials of higher order. We also define new generating functions of multiple q -Euler numbers and polynomials. Furthermore, we construct Euler q -Zeta function.

1. INTRODUCTION

For any complex number z , it is well known that the familiar Euler polynomials $E_n(z)$ are defined by means of the following generating function, cf.[3, 5, 6, 9, 13]:

$$F(z, t) = \frac{2}{e^t + 1} e^{zt} = \sum_{n=0}^{\infty} E_n(z) \frac{t^n}{n!}, \quad (|t| < \pi). \quad (1)$$

We note that, by substituting $z = 0$ into (1), $E_n(0) = E_n$ is the familiar n -th Euler number defined by

$$G(t) = F(0, t) = \frac{2}{e^t + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \quad (|t| < \pi), \text{ cf. [4, 5].}$$

By the meaning of the generalization of E_n , Frobenius-Euler numbers and polynomials are also defined by

$$\frac{1-u}{e^t - u} = \sum_{n=0}^{\infty} H_n(u) \frac{t^n}{n!}, \text{ and } \frac{1-u}{e^t - u} e^{xt} = \sum_{n=0}^{\infty} H_n(u, x) \frac{t^n}{n!} \quad (u \in \mathbb{C} \text{ with } |u| > 1), \text{ cf. [16].}$$

Over five decades ago, Calitz [2, 3] defined q -extension of Frobenius-Euler numbers and polynomials and proved properties analogous to those satisfied $H_n(u)$ and $H_n(u, x)$. Recently, Satoh [14, 15] used these properties, especially the so-called distribution relation for the q -Frobenius-Euler polynomials, in order to construct the corresponding q -extension of the p -adic measure and to define a q -extension of p -adic l -function $l_{p,q}(s, u)$.

Let p be a fixed odd prime in this paper. Throughout this paper, the symbols $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$, denote the ring of rational integers, the ring of p -adic integers, the field of p -adic numbers, the complex number field, and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let $\nu_p(p)$ be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one speaks of q -extension, q can be regarded as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$; it is always clear from the context. If $q \in \mathbb{C}$, then one usually assumes that $|q| < 1$. If $q \in \mathbb{C}_p$, then one usually assumes that $|q - 1|_p < p^{-\frac{1}{p-1}}$, and hence $q^x = \exp(x \log q)$ for $x \in \mathbb{Z}_p$. In this paper, we use the below notation

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad (a : q)_n = (1 - a)(1 - aq) \cdots (1 - aq^{n-1}), \text{ cf. [6, 7, 8, 9, 10, 11, 14] }.$$

Note that $\lim_{q \rightarrow 1} [x]_q = x$ for any x with $|x|_p \leq 1$ in the p -adic case. For a fixed positive integer d with $(p, d) = 1$, set

$$\begin{aligned} X &= X_d = \varprojlim_N \mathbb{Z}/dp^N, \\ X_1 &= \mathbb{Z}_p, X^* = \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dp\mathbb{Z}_p, \\ a + dp^N\mathbb{Z}_p &= \{x \in X | x \equiv a \pmod{p^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^N$, (cf. [10, 11]). We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$, and write $f \in UD(\mathbb{Z}_p)$, if the difference quotients $F_f(x, y) = \frac{f(x) - f(y)}{x - y}$ have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$, cf. [11]. For $f \in UD(\mathbb{Z}_p)$, let us begin with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{0 \leq j < p^N} f(j) \mu_q(j + p^N\mathbb{Z}_p), \text{ cf. [7, 8, 9, 11],}$$

which represents a q -analogue of Riemann sums for f . The integral of f on $\mathbb{Z}_p$ is defined as the limit of those sums (as $n \rightarrow \infty$) if this limit exists. The q -Volkenborn integral of a function $f \in UD(\mathbb{Z}_p)$ is defined by

$$I_q(f) = \int_X f(x) d\mu_q(x) = \int_{X_d} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[dp^N]_q} \sum_{x=0}^{dp^N-1} f(x) q^x.$$

Recently, we considered another construction of a q -Eulerian numbers, which are different than Carlitz's q -Eulerian numbers as follows, cf.[6, 12, 13]:

$$F_q(x, t) = [2]_q \sum_{n=0}^{\infty} (-1)^n q^n e^{[n+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}.$$

Thus we have

$$E_{n,q} = E_{n,q}(0) = \frac{[2]_q}{(1-q)_n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}},$$

$$E_{n,q}(x) = \frac{[2]_q}{(1-q)_n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}} q^{lx},$$

where $\binom{n}{l}$ is a binomial coefficient, cf.[13].

Note that $\lim_{q \rightarrow 1} E_{n,q} = E_n$ and $\lim_{q \rightarrow 1} E_{n,q}(x) = E_n(x)$. In [12], we also proved that q -Eulerian polynomial $E_{n,q}(x)$ can be represented by q -Volkenborn integral as follows:

$$\int_{X_d} [x + x_1]_q^k d\mu_{-q}(x_1) = \int_{\mathbb{Z}_p} [x + x_1]_q^k d\mu_{-q}(x) = E_{k,q}(x), \quad \text{for } k, d \in \mathbb{N},$$

where $\mu_{-q}(x + p^N \mathbb{Z}_p) = \frac{q^x [2]_q}{1+q^{p^N}} (-1)^x$.

The purpose of this paper is to present a systemic study of some families of multiple q -Euler numbers and polynomials. In particular, by using the q -Volkenborn integration on $\mathbb{Z}_p$, we construct p -adic q -Euler numbers and polynomials of higher order. We also define new generating function of these q -Euler numbers and polynomials of higher order. Furthermore, we construct Euler q - ζ -function. From section 2 to section 5, we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-\frac{1}{p-1}}$.

2. q -EULER NUMBERS AND POLYNOMIALS ASSOCIATED WITH AN INVARIANT p -ADIC q -INTEGRALS ON $\mathbb{Z}_p$

Let $h \in \mathbb{Z}$, $k \in \mathbb{N} = \{1, 2, 3, \dots\}$, and let us consider the extended higher-order q -Euler numbers as follows:

$$E_{m,q}^{(h,k)} = \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x_1 + x_2 + \cdots + x_k]_q^m q^{x_1(h-1) + \cdots + x_k(h-k)} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k).$$

Then we have

$$E_{m,q}^{(h,k)} = \frac{[2]_q^k}{(1-q)^m} \sum_{l=0}^m \binom{m}{l} \frac{(-1)^l}{(-q^{h+l} : q^{-1})_k}.$$

From the definition of $E_{m,q}^{(h,k)}$, we can easily derive the below:

$$E_{m,q}^{(h,k)} = E_{m,q}^{(h-1,k)} + (q-1)E_{m+1,q}^{(h-1,k)}, \quad (m \geq 0).$$

It is easy to show that

$$\begin{aligned} & \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k+1 \text{ times}} q^{\sum_{j=1}^{k+1} (m-j)x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_{k+1}) \\ &= \sum_{j=1}^m \binom{m}{j} (q-1)^j \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k+1 \text{ times}} \left[\sum_{l=1}^{k+1} x_l \right]_q^j q^{-\sum_{j=1}^k j x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_{k+1}), \end{aligned} \quad (2)$$

and we also get

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k+1 \text{ times}} q^{\sum_{j=1}^{k+1} (m-j)x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_{k+1}) = \frac{[2]_q^{k+1}}{(-q^m : q^{-1})_{k+1}}. \quad (3)$$

From (2) and (3), we can derive the below proposition.

Proposition 1. *For $m, k \in \mathbb{N}$, we have*

$$\begin{aligned} & \sum_{j=0}^m \binom{m}{j} (q-1)^j E_{j,q}^{(0,k+1)} = \frac{[2]_q^{k+1}}{(-q^m : q^{-1})_{k+1}}, \\ & E_{m,q}^{(h,k)} = \frac{[2]_q^k}{(1-q)^m} \sum_{l=0}^m \binom{m}{l} \frac{(-1)^l}{(-q^{h+l} : q^{-1})_k}. \end{aligned}$$

Remark. Note that $E_{n,q}^{(1,1)} = E_{n,q}$, where $E_{n,q}$ are the q -Euler numbers (see [13]).

From the definition of $E_{n,q}^{(h,k)}$, we can derive

$$\sum_{j=0}^i \binom{i}{j} (q-1)^j E_{m-i+j,q}^{(h-1,k)} = \sum_{j=0}^{i-1} \binom{i-1}{j} (q-1)^j E_{m+j-i,q}^{(h,k)}$$

for $m \geq i$. By simple calculation, we easily see that

$$\sum_{j=0}^m \binom{m}{j} (q-1)^j E_{j,q}^{(h,1)} = \int_{\mathbb{Z}_p} q^{mx} q^{(h-1)x} d\mu_{-q}(x) = \frac{[2]_q}{[2]_q^{m+h}}.$$

Furthermore, we can give the following relation for the q -Euler numbers, $E_{m,q}^{(0,h)}$;

$$\sum_{j=0}^m \binom{m}{j} (q-1)^j E_{j,q}^{(0,k)} = \frac{[2]_q^k}{(-q^m : q^{-1})_k}. \quad (4)$$

3. POLYNOMIALS $E_{n,q}^{(0,k)}(x)$

We now define the polynomials $E_{n,q}^{(0,k)}(x)$ (in q^x) by

$$E_{n,q}^{(0,k)}(x) = \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x_1 + x_2 + \cdots + x_k]_q^m q^{\sum_{j=1}^k j x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k).$$

Thus, we have

$$(q-1)^m E_{m,q}^{(0,k)}(x) = [2]_q^k \sum_{j=0}^m \binom{m}{j} q^{jx} (-1)^{m-j} \frac{1}{(-q^j : q^{-1})_k}. \quad (5)$$

It is not difficult to show that

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} q^{\sum_{j=1}^m (m-j)x_j + mx} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) = q^{mx} \frac{[2]_q^k}{(-q^m : q^{-1})_k},$$

and

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} q^{\sum_{j=1}^m (m-j)x_j + mx} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) = \sum_{j=0}^m \binom{m}{j} (q-1)^j E_{j,q}^{(0,k)}(x).$$

Therefore we obtain the following.

Lemma 2. For $m, k \in \mathbb{N}$, we have

$$\begin{aligned} \sum_{j=0}^m \binom{m}{j} (q-1)^j E_{j,q}^{(0,k)}(x) &= \frac{q^{mx} [2]_q^k}{(-q^m : q^{-1})_k}, \\ E_{m,q}^{(0,k)}(x) &= \frac{[2]_q^k}{(1-q)^m} \sum_{j=0}^m \binom{m}{j} q^{jx} (-1)^j \frac{1}{(-q^j : q^{-1})_k}. \end{aligned} \quad (6)$$

Let $l \in \mathbb{N}$ with $l \equiv 1 \pmod{2}$. Then we get easily

$$\begin{aligned} &\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} \left[x + \sum_{j=1}^k x_j \right]_q^m q^{-\sum_{j=1}^k j x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \frac{[l]_q^m}{[l]_{-q}^k} \sum_{i_1, \dots, i_k=0}^{l-1} q^{-\sum_{j=2}^k (j-1)i_j} \\ &\quad \cdot (-1)^{\sum_{j=1}^k i_j} \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} \left[\frac{x + \sum_{j=1}^k i_j}{l} + \sum_{j=1}^k x_j \right]_{q^l}^m q^{-l \sum_{j=1}^k j x_j} d\mu_{-q^l}(x_1) \cdots d\mu_{-q^l}(x_k). \end{aligned}$$

From this, we can derive the following “multiplication formula”:

Theorem 3. *Let l be an odd positive integer. Then*

$$E_{m,q}^{(0,k)}(x) = \frac{[l]_q^m}{[l]_{-q}^k} \sum_{i_1, \dots, i_k=0}^{l-1} q^{-\sum_{j=2}^k (j-1)i_j} (-1)^{\sum_{i=1}^k i_l} E_{m,q^l}^{(0,k)}\left(\frac{x + i_1 + \dots + i_k}{l}\right). \quad (7)$$

Moreover,

$$E_{m,q}^{(0,k)}(lx) = \frac{[l]_q^m}{[l]_{-q}^k} \sum_{i_1, \dots, i_k=0}^{l-1} q^{-\sum_{j=2}^k (j-1)i_j} (-1)^{\sum_{i=1}^k i_l} E_{m,q^l}^{(0,k)}\left(x + \frac{i_1 + \dots + i_k}{l}\right). \quad (8)$$

From (4) and (5), we can also derive the below expression for $E_{n,q}^{(0,k)}(x)$:

$$E_{m,q}^{(0,k)}(x) = \sum_{i=0}^m \binom{m}{i} E_{i,q}^{(0,k)}[x]_q^{m-i} q^{ix}, \quad (9)$$

whence also

$$E_{m,q}^{(0,k)}(x+y) = \sum_{j=0}^m \binom{m}{j} [y]_q^{m-j} q^{jy} E_{j,q}^{(0,k)}(x). \quad (10)$$

4. POLYNOMIALS $E_{m,q}^{(h,1)}(x)$

Let us define

$$E_{m,q}^{(h,1)}(x) = \int_{\mathbb{Z}_p} [x + x_1]_q^m q^{x_1(h-1)} d\mu_{-q}(x_1). \quad (11)$$

Then we have

$$E_{m,q}^{(h,1)}(x) = \frac{[2]_q}{(1-q)^m} \sum_{l=0}^m \binom{m}{l} (-1)^l q^{lx} \frac{1}{(1+q^{l+h})}.$$

By simple calculation of q -Volkenvorn integral, we note that

$$\begin{aligned} q^x \int_{\mathbb{Z}_p} [x + x_1]_q^m q^{x_1(h-1)} d\mu_{-q}(x_1) &= (q-1) \int_{\mathbb{Z}_p} [x + x_1]_q^{m+1} q^{x_1(h-2)} d\mu_{-q}(x_1) \\ &\quad + \int_{\mathbb{Z}_p} [x + x_1]_q^m q^{x_1(h-2)} d\mu_{-q}(x_1). \end{aligned}$$

Thus, we have

$$q^x E_{m,q}^{(h,1)}(x) = (q-1) E_{m+1,q}^{(h-1,1)}(x) + E_{m,q}^{(h-1,1)}(x). \quad (12)$$

It is easy to show that

$$\int_{\mathbb{Z}_p} [x + x_1]_q^m q^{(h-1)x_1} d\mu_{-q}(x_1) = \sum_{j=0}^m \binom{m}{j} [x]_q^{m-j} q^{jx} \int_{\mathbb{Z}_p} [x_1]_q^j q^{(h-1)x_1} d\mu_{-q}(x_1).$$

This is equivalent to

$$\begin{aligned} E_{m,q}^{(h,1)}(x) &= \sum_{j=0}^m \binom{m}{j} [x]_q^{m-j} q^{jx} E_{j,q}^{(h,1)} \\ &= \left(q^x E_q^{(h,1)} + [x]_q \right)^m, \quad \text{for } m \geq 1, \end{aligned}$$

where we use the technique method notation by replacing $(E_q^{(h,1)})^n$ by $E_{n,q}^{(h,1)}$, symbolically. From (11), we can derive

$$q^h E_{m,q}^{(h,1)}(x+1) + E_{m,q}^{(h,1)}(x) = [2]_q [x]_q^m. \quad (13)$$

For $x = 0$ in (13), this gives

$$q^h \left(q E_{m,q}^{(h,1)} + 1 \right)^m + E_{m,q}^{(h,1)} = \delta_{0,k}, \quad (14)$$

where $\delta_{0,k}$ is Kronecker symbol. By the simple calculation of q -Volkenborn integration, we easily see that

$$\int_{\mathbb{Z}_p} q^{x_1(h-1)} d\mu_{-q}(x_1) = \frac{[2]_q}{[2]_{q^h}}.$$

Thus, we have $E_{0,q}^{(h,1)} = \frac{[2]_q}{[2]_{q^h}}$. From the definition of q -Euler polynomials, we can derive

$$\int_{\mathbb{Z}_p} [1 - x + x_1]_{q-1}^m q^{-x_1(h-1)} d\mu_{-q}(x_1) = q^{m+h-1} (-1)^m E_{m,q}^{(h,1)}(x).$$

Therefore we obtain the below “complementary formula”:

Theorem 4. For $m \in \mathbb{N}$, $n \in \mathbb{Z}$, we have

$$E_{m,q^{-1}}^{(h,1)}(1-x) = (-1)^m q^{m+h-1} E_{m,q}^{(h,1)}(x). \quad (15)$$

In particular, for $x = 1$, we see that

$$\begin{aligned} E_{m,q^{-1}}^{(h,1)}(0) &= (-1)^m q^{m+h-1} E_{m,q}^{(h,1)}(1) \\ &= (-1)^{m-1} q^{m-1} E_{m,q}^{(h,1)}, \quad \text{for } m \geq 1. \end{aligned} \quad (16)$$

For $l \in \mathbb{N}$ with $l \equiv 1 \pmod{2}$, we have

$$\begin{aligned} &\int_{\mathbb{Z}_p} q^{(h-1)x_1} [x + x_1]_q^m q^{x_1(h-1)} d\mu_{-q}(x_1) \\ &= \frac{[l]_q^m}{[l]_{-q}} \sum_{i=0}^{l-1} q^{hi} (-1)^i \int_{\mathbb{Z}_p} \left[\frac{x+i}{l} + x_1 \right]_{q^l}^m q^{x_1(h-1)l} d\mu_{-q^l}(x_1). \end{aligned}$$

Thus, we can also obtain the following:

Theorem 5. (*Multiplication formula*) For $l \in \mathbb{N}$ with $l \equiv 1 \pmod{2}$, we have

$$\frac{[2]_q}{[2]_{q^l}} [l]_q^m \sum_{i=0}^{l-1} q^{hi} (-1)^i E_{m,q^l}^{(h,1)}\left(\frac{x+i}{l}\right) = E_{m,q}^{(h,1)}(x).$$

Furthermore,

$$\frac{[2]_q}{[2]_{q^l}} [l]_q^m \sum_{i=0}^{l-1} q^{hi} (-1)^i E_{m,q^l}^{(h,1)}\left(x + \frac{i}{l}\right) = E_{m,q}^{(h,1)}(lx).$$

5. POLYNOMIALS $E_{m,q}^{(h,k)}(x)$ AND $h = k$

It is now easy to combine the above results and define the new polynomials as follows:

$$E_{m,q}^{(h,k)}(x) = \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x + x_1 + \cdots + x_k]_q^m q^{(h-1)x_1 + \cdots + (h-k)x_k} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k).$$

Thus, we note that

$$(q-1)^m E_{m,q}^{(h,k)}(x) = \sum_{j=0}^m \binom{m}{j} (-1)^{m-j} q^{xj} \frac{[2]_q^k}{(-q^{j+h} : q^{-1})_k}. \quad (17)$$

We may now mention the following formulas which are easy to prove.

$$q^h E_{m,q}^{(h,k)}(x+1) + E_{m,q}^{(h,k)}(x) = [2]_q E_{m,q}^{(h-1,k-1)}(x), \quad (18)$$

and

$$q^x E_{m,q}^{(h+1,k)}(x) = (q-1) E_{m+1,q}^{(h,k)}(x) + E_{m,q}^{(h,k)}(x). \quad (19)$$

Let $l \in \mathbb{N}$ with $l \equiv 1 \pmod{2}$. Then we note that

$$\begin{aligned} & \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} \left[x + \sum_{j=1}^k x_j \right]_q^m q^{\sum_{j=1}^k (h-j)x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \frac{[l]_q^m}{[l]_{-q}^k} \sum_{i_1, \dots, i_k=0}^{l-1} q^{h \sum_{j=1}^k i_j - \sum_{j=2}^k (j-1)i_j} (-1)^{\sum_{j=1}^k i_j} \\ & \cdot \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} \left[\frac{x + \sum_{j=1}^k i_j}{l} + \sum_{j=1}^k x_j \right]_{q^l}^m (q^l)^{\sum_{j=1}^k (h-j)x_j} d\mu_{-q^l}(x_1) \cdots d\mu_{-q^l}(x_k). \end{aligned}$$

Therefore we obtain the following:

Theorem 6. (*Distribution for q -Euler polynomials*) For $l \in \mathbb{N}$ with $l \equiv 1 \pmod{2}$. Then we have

$$(20) \quad E_{m,q}^{(h,k)}(lx) = \frac{[l]_q^m}{[l]_q^{-q}} \sum_{i_1, \dots, i_k=0}^{l-1} q^{h \sum_{j=1}^k i_j - \sum_{j=2}^k (j-1)i_j} (-1)^{\sum_{j=1}^k i_j} E_{m,q^l}^{(h,k)} \left(x + \frac{i_1 + \dots + i_k}{l} \right).$$

It is interesting to consider the case $h = k$, which leads to the desired extension of the q -Euler numbers of higher order, cf.[1]. We shall denote the polynomials in this special case by $E_{m,q}^{(k)}(x) := E_{m,q}^{(k,k)}(x)$. Then we have

$$(q-1)^m E_{m,q}^{(k)}(x) = \sum_{j=0}^m \binom{m}{j} (-1)^{m-j} q^{jx} \frac{[2]_q^k}{(-q^{j+k} : q^{-1})_k}, \quad (21)$$

and

$$E_{m,q^{-1}}^{(k)}(k-x) = (-1)^m q^{m+\binom{k}{2}} E_{m,q}^{(k)}(x). \quad (22)$$

For $x = k$ in (22), we see that

$$E_{m,q^{-1}}^{(k)}(0) = (-1)^m q^{m+\binom{k}{2}} E_{m,q}^{(k)}(k). \quad (23)$$

From (18), we can derive the below formula:

$$q^k E_{m,q}^{(k)}(x+1) + E_{m,q}^{(k)}(x) = [2]_q E_{m,q}^{(k-1)}(x). \quad (24)$$

Putting $x = 0$ in (17), we obtain

$$(q-1)^m E_{m,q}^{(k)} = \sum_{i=0}^m \binom{m}{i} (-1)^{m-i} \frac{[2]_q^k}{(-q^{i+k} : q^{-1})_k}. \quad (25)$$

Note that

$$\begin{aligned} & \sum_{i=0}^m \binom{m}{i} (q-1)^i \underbrace{\int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x_1 + \dots + x_k]_q^i q^{\sum_{j=1}^{k-1} (k-j)x_j} d\mu_{-q}(x_1) \dots d\mu_{-q}(x_k) \\ &= \frac{[2]_q^k}{(-q^{m+k} : q^{-1})_k}. \end{aligned}$$

From this, we can easily derive

$$\sum_{i=0}^m \binom{m}{i} (q-1)^i E_{i,q}^{(k)} = \frac{[2]_q^k}{(-q^{m+k} : q^{-1})_k} \quad (26)$$

and so it follows

$$E_{m,q}^{(k)}(x) = (q^x E_q^{(k)} + [x]_q)^m, \quad m \geq 1, \quad (27)$$

where we use the technique method notation by replacing $(E_q^{(k)})^n$ by $E_{n,q}^{(k)}$, symbolically. In particular, from (24), we have

$$q^k(qE_q^{(k)} + 1)^m + E_{m,q}^{(k)} = [2]_q E_{m,q}^{(k-1)}. \quad (28)$$

It is easy to see that

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} q^{(k-1)x_1 + \cdots + x_{k-1}} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) = \frac{[2]_q^k}{(-q^k : q^{-1})_k}.$$

Thus, we note that $E_{0,q}^{(k)} = \frac{[2]_q^k}{(-q^k : q^{-1})_k}$.

6. GENERATING FUNCTION FOR q -EULER POLYNOMIALS

An obvious generating function for q -Euler polynomials is obtained, from (17), by

$$\begin{aligned} & [2]_q^k e^{\frac{t}{1-q}} \sum_{j=0}^{\infty} \frac{(-1)^j}{(-q^{j+h} : q^{-1})_k} q^{jx} \left(\frac{1}{1-q} \right)^j \frac{t^j}{j!} \\ &= \sum_{n=0}^{\infty} E_{n,q}^{(h,k)} \frac{t^n}{n!}. \end{aligned} \quad (29)$$

From (17), we can also derive the below formula:

$$q^{h-k} E_{m,q}^{(h,k+1)}(x+1) = [2]_q E_{m,q}^{(h,k)}(x) - E_{m,q}^{(h,k+1)}(x). \quad (30)$$

Again from (21) and (25), we get easily

$$\begin{aligned} & \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} \left[x + \sum_{j=1}^k x_j \right]_q^m q^{\sum_{j=1}^{k-1} (k-j)x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \sum_{j=0}^m \binom{m}{j} q^{xj} \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x_k]_q^j \left[x + \sum_{j=1}^{k-1} x_j \right]_q^{n-j} q^{\sum_{l=1}^{k-1} (k+j-l)x_l} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k). \end{aligned}$$

Thus, we note that

$$E_{m,q}^{(k)}(x) = \sum_{j=0}^m \binom{m}{j} q^{xj} E_{j,q}^{(1)} E_{m-j,q}^{(k+j,k-1)}(x). \quad (31)$$

Take $x = 0$ in (31), we have

$$E_{m,q}^{(k)} = \sum_{i=0}^m \binom{m}{i} E_{i,q}^{(1)} E_{m-i,q}^{(k+i,k-1)}. \quad (32)$$

So, for $k = 2$,

$$E_{m,q}^{(2)} = \sum_{i=0}^m \binom{m}{i} E_{i,q} E_{m-i,q}^{(j+2,1)}.$$

It is not difficult to show that

$$\int_{\mathbb{Z}_p} [x]_q^m q^{hx} d\mu_{-q}(x) = \sum_{j=0}^h \binom{h}{j} (q-1)^j \int_{\mathbb{Z}_p} [x]_q^{m+j} d\mu_{-q}(x), \text{ for } h \in \mathbb{N}.$$

From this, we can derive the below:

$$E_{m,q}^{(h+1,1)} = \sum_{j=0}^h \binom{h}{j} (q-1)^j E_{m+j,q}, \quad h \in \mathbb{N}. \quad (33)$$

By (32) and (33), we easily see that

$$E_{m,q}^{(2)} = \sum_{j=0}^m \binom{m}{j} E_{j,q} \sum_{i=0}^{j+1} \binom{j+1}{i} (q-1)^i E_{m-j+i,q}. \quad (34)$$

By (34), for $q = 1$, we note that

$$E_m^{(2)} = \sum_{j=0}^m \binom{m}{j} E_j E_{m-j}, \text{ where } \left(\frac{2}{e^t + 1} \right)^k = \sum_{n=0}^{\infty} E_n^{(k)} \frac{t^n}{n!}.$$

It is easy to show that

$$[x + x_1 + \cdots + x_k]_q^m = \sum_{j=0}^m \binom{m}{j} [x_1 + x]_q^{m-j} q^{j(x_1+x)} [x_2 + \cdots + x_k]_q^j.$$

By using this, we get easily

$$\begin{aligned}
& \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x + \sum_{j=1}^k x_j]_q^m q^{\sum_{j=1}^{k-1} (k-j)x_j} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\
&= \sum_{j=0}^m \binom{m}{j} q^{jx} \int_{\mathbb{Z}_p} [x + x_1]_q^{m-j} q^{(k+j-1)x_1} d\mu_{-q}(x_1) \\
&\quad \cdot \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k-1 \text{ times}} [x_2 + \cdots + x_k]_q^j q^{\sum_{j=2}^{k-1} (k-j)x_j} d\mu_{-q}(x_2) \cdots d\mu_{-q}(x_k).
\end{aligned}$$

Therefore we obtain the following:

Theorem 7. *For $m, k \in \mathbb{N}$, we have*

$$E_{m,q}^{(k)}(x) = \sum_{j=0}^m \binom{m}{j} q^{jx} E_{m-j,q}^{(k+j,1)}(x) E_{j,q}^{(k-1)}. \quad (35)$$

Indeed for $x = 0$,

$$\begin{aligned}
E_{m,q}^{(k)} &= \sum_{j=0}^m \binom{m}{j} E_{m-j,q}^{(k+j,1)} E_{j,q}^{(k-1)} \\
&= \sum_{j=0}^m \binom{m}{j} E_{j,q}^{(k-1)} \sum_{i=0}^{k+j} (q-1)^i \binom{k+j-1}{i} E_{m-j+i,q}^{(1)}.
\end{aligned} \quad (36)$$

As for $q = 1$, we get the below formula

$$E_m^{(k)} = \sum_{j=0}^m \binom{m}{j} E_j^{(k-1)} E_{m-j}^{(1)}.$$

7. q -EULER ZETA FUNCTION IN $\mathbb{C}$

In this section, we assume that $q \in \mathbb{C}$ with $|q| < 1$. From section 4, we note that

$$\begin{aligned}
E_{m,q}^{(h,1)}(x) &= \frac{[2]_q}{(1-q)^m} \sum_{l=0}^m \binom{m}{l} q^{lx} (-1)^l \frac{1}{1+q^{l+h}} \\
&= [2]_q \sum_{n=0}^{\infty} (-1)^n q^{nh} [n+x]_q^n.
\end{aligned} \quad (37)$$

Thus, we can define q -Euler zeta function:

Defintion 8. For $s, q \in \mathbb{C}$ with $|q| < 1$, define

$$\zeta_{E,q}^h(s, x) = [2]_q \sum_{n=0}^{\infty} \frac{(-1)^n q^{nh}}{[n+x]_q^s},$$

where $x \in \mathbb{R}$ with $0 < x \leq 1$.

Note that $\zeta_{E,q}^h(-m, x) = E_{m,q}^{(h,1)}(x)$, for $m \in \mathbb{N}$. Let

$$F_q(t, x) = \sum_{n=0}^{\infty} E_{n,q}^{(h,1)}(x) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned} F_q(t, x) &= [2]_q e^{\frac{t}{1-q}} \sum_{n=0}^{\infty} (-1)^n q^{hn} e^{-\frac{q^{n+x}}{1-q} t} \\ &= [2]_q \sum_{n=0}^{\infty} (-1)^n q^{hn} e^{[n+x]_q t}, \quad \text{for } h \in \mathbb{Z}. \end{aligned}$$

Therefore we obtain the following

Lemma 9. For $h \in \mathbb{Z}$, we have

$$\begin{aligned} F_q(t, x) &= [2]_q \sum_{n=0}^{\infty} (-1)^n q^{hn} e^{[n+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,q}^{(h,1)}(x) \frac{t^n}{n!}. \end{aligned} \tag{38}$$

Let $\Gamma(s)$ be the gamma function. Then we easily see that

$$\frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-1} F_q(-t, x) dt = \zeta_{E,q}^h(s, x), \quad \text{for } s \in \mathbb{C}. \tag{39}$$

From (38) and (39), we can also derive the below Eq.(40):

$$(40) \quad \zeta_{E,q}^h(-n, x) = E_{n,q}^{(h,1)}(x), \quad \text{for } n \in \mathbb{N}.$$

REFERENCES

- [1] E. Boros, K. Elbassioni, V. Gurvich, L. Khachiyan, K. Makino, *An intersection inequality for discrete distributions and related generation problems*, Lecture Notes in Comput. Sci. **2719** (2003), 543-555.
- [2] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
- [3] L. Carlitz, *q-Bernoulli and Eulerian numbers*, Trans. Amer. Math. Soc. **76** (1954), 332-350.
- [4] E. Deeba and D. Rodriguez,, *Stirling's series and Bernoulli numbers*, Amer. Math. Monthly **98** (1991), 423-426.
- [5] T. Howard, *Applications of a recurrences formula for the Bernoulli numbers*, J. Number Theory **52** (1995), 157-172.
- [6] T. Kim, L. C. Jang, H. K. Park, *A note on q-Euler and Genocchi numbers*, Proc. Japan Academy **77** (2001), 139-141.
- [7] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12** (2005), 189-196.
- [8] T. Kim, *Non-Archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials*, Russ. J. Math. Phys. **10** (2003), 91-98.
- [9] T. Kim, *An invariant p-adic integral associated with Daehee numbers*, Integral Trnasforms and special functions **13** (2002), 65-69.
- [10] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320-329.
- [11] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [12] T. Kim, *A note on q-Volkenborn integration*, Proc. Jangjeon Math. Soc. **8** (2005), 13-17.
- [13] T. Kim, *q-Euler and Genocchi numbers*, arXiv: math. NT/0506278 vol 1 14 June (2005).
- [14] J. Satoh, *q-analogue of Riemann's ζ -function and q-Euler numbers*, J. Number Theory **31** (1989), 346-362.
- [15] J. Satoh, *Sums products of two q-Bernoulli numbers*, J. Number Theory **74** (1999), 173-180.
- [16] J. Shiratani and S. Yamamoto, *On a p-adic interpolating function for the Euler numbers and its derivatives*, Mem. Fac. Sci. Kyushu Univ. Math. **39** (1985), 113-125.